

Facilitators for Clouds critique? Friday, Monday and Weds?

Max
Matthew Mc



• Cloud image submission: Include

- 1) your edited image
- 2) your original (unedited) image
- 3) the appropriate Skew-T diagram
- 4) a short statement of cloud type and stable or unstable atm.
- 5) Post on Flowvis.org. Edit your post date to match your cloud date and time.

Post

Canvas Tonight

in report

in report after critique

FLOWVIS.ORG

Clouds = droplets or ice MOVING UPWARDS

Lift mechanisms:

1. Instability: creates Cumulus clouds
2. Orographics: terrain, mountains
3. Synoptic scale weather systems; local instability. Both at warm and cold fronts; cold air pushes under in a cold front, warm air overruns in a warm front.
4. Convergence: shoreline temperature differences and cyclonic uplift

finish up

1. Instability driven clouds

Flat / Puffy

Stratocumulus: probably the world's most common cloud.

Stratocumulus
Formation mechanisms:

<http://www.flowvis.org/category/flow-categories/clouds/stratocumulus/>




- 1) Cumulus joined together, caused by an inversion, a stable layer that stops upward convection
- 2) Stratus broken up. Top reflects UV, visible light, cools (maybe radiates IR to space). Bottom absorbs IR from the earth, warms. Cool on top, warm on the bottom = unstable, wants to turn over, breaking up stratus layer. Stratocumulus stratiformus



<http://www.flowvis.org/2013/04/11/stratocumulus-boulder-co-18th-of-february-2013-at-1131-a-m/>

Bottom absorbs IR from the earth, warms
Cool on top, warm on the bottom = unstable,
wants to turn over, breaking up stratus layer.
Stratocumulus stratiformis



instability

Partial rule of thumb
Cumulus = from instability; local uplift
Stratus = more stable, from widespread uplift

These are GENUS

For info on Species, Varieties and Accessory Clouds, see

Interesting book on how clouds were first classified and named ~1804, by Luke Howard
Richard Hamblin, *The Invention of Clouds: How an Amateur Meteorologist Forged the Language of the Skies* (Picador, 2002).

Another rule of thumb (fingers, really)
Measure cloud element size with hand outstretched.
Cirrocumulus = elements smaller than one finger width
Alto cumulus = elements between one and three finger widths
Cumulus = elements larger than three finger widths.

Also finish up

2: Orographic clouds, caused by topography, i.e. mountains
Orography (from the Greek ὄρος, hill, γράφω, to write) [Wikipedia]

Most common interesting cloud in winter and spring is the


Alto cumulus ^{standing} lenticularis (higher than 6500 ft above local ground level) ACSL
or
Stratocumulus lenticularis (lower)
or
Mountain Wave Cloud, trapped or lee

requires STABLE atmosphere: note exception to unstable/cumulus pairing

STANDING WAVE

Clouds Produced by Vertically **Trapped** Mountain Waves

Thomas Carney et al.,
AC 00-57 Hazardous
Mountain Winds and
Their Visual Indicators
(Federal Aviation
Administration, 1997),
http://rgl.faa.gov/Regulatory_and_Guidance_Library/rgAdvisoryCircular.nsf/0/780437D88CBDAFD086256A94006FD5B8?OpenDocument.




Clouds that sit right on the Divide =
FOEHN cloud wall.
From air being forced up over the mountains

Fayne

Alto cumulus lenticularis. Typically 1 to 5 wave crests.

Clouds stay stationary, but may move off and reform periodically



AGSL

Foehn

1-5 wave crest

Ben Britton, FV 2010

it covers sky

If there's more wave crests, or short wavelengths, it's probably NOT a mountain wave cloud; more likely altocumulus undulatus, from gravity waves in the atmosphere, like ripples on a liquid surface.

<http://www.colorado.edu/MCEN/flowvis/galleries/2007/assignment2.html>



from synoptic lift



1141


Tracy Eliasson FV 2007

Could also be from wind shear, via the Kelvin Helmholtz instability



Rare to be able to see cross section like this

<http://cloudappreciationsociety.org/collecting/terry-robinson/>




Minute paper: Which way is the wind going?
Where is it faster?



Colin Stewart FV 2012 Clouds 1




all static





3: Synoptic uplift = weather system clouds.

Weather system progressions; 'synoptic scale' uplifts (1000 km across).
Any type of cloud is possible.



Inserted from: <file:///C:/Users/hertzber1/Documents/01CLASSES/FlowVis/Content/scanned images/TypWeatherSystem.tif>





Low Pressure System: Air tries to move into low, Coriolis makes it turn left = counterclockwise circulation. Typically unstable. **CYCLONIC**
CCW

High pressure system: Air tries to move out. Coriolis makes it turn right = clockwise circulation. Weak or nonexistent fronts, so no instability. **ANTICYCLONIC**



Idealized depiction of the support that divergence and convergence aloft provide to cyclonic and anticyclonic circulation at the surface.

<http://earth.usc.edu>




Idealized depiction of the support that divergence and convergence aloft provide to cyclonic and anticyclonic circulation at the surface.

<http://earth.usc.edu/~stott/Catalina/WeatherPatterns.html>

Divergence aloft creates convergence and lift at surface. Pumping action. Bad for wildland fires.

4: Convergence uplift along shorelines



Cool sea breeze is pulled in during daytime. **onshore**
Land or shore breeze happens at night, when land cools more rapidly than the water.
Note: winds are named for where they come from

CloudClassificationTable.pdf; Copyrighted, but available in D2L.
Also see [Cloud types for observers \(PDF, 4 MB\) - Met Office](#) 45 pgs

The Cloud Spotter's Guide
 CLOUD CLASSIFICATION TABLE
 Gayin Pretor-Pinney, Berigee Press 2006
 Clouds are classified according to a Latin "Linnaean" system (similar to the one used for plants and animals), which is based on their heights and appearance. Most clouds fall into one of ten basic groups, known as 'genera'. They can further be defined as one of the possible 'species' for that genus, and any combination of the possible 'varieties'. There are also various accessory clouds and supplementary features that sometimes appear in conjunction with the main cloud types. (If all this Latin freaks you out, don't worry - it freaks me out too.)

GENUS	SPECIES (CAN ONLY BE ONE)	VARIETIES (CAN BE MORE THAN ONE)	ACCESSORY CLOUDS AND SUPPLEMENTARY FEATURES	
Circulus	humilis		pernis	arcs
	mediocris	radiatus	velum	parvus
	congestus		virga	tuba
	fractus		precipitatio	
Cumulonimbus (forms through all three levels)	calvus	(none)	virga	velum
	capillatus	(none)	pernis	arcs
			lucos	tuba
Stratus	nebulosus	opacus		
	fractus	translucidus	precipitatio	
		undulatus		
Stratocumulus		translucidus		
		perfractus		
	stratiformis	opacus	mamma	
	lenticularis	duplicatus	virga	
	castellatus	undulatus	precipitatio	
		radiatus		
Altostratus		lacunosus		
		translucidus		
	stratiformis	perfractus		
	lenticularis	opacus	virga	
	castellatus	duplicatus	mamma	
	floccus	undulatus		
Altostratus		radiatus	virga	
		opacus	precipitatio	
	(none)	duplicatus	pernis	
		undulatus	mamma	
Nimbostratus (forms through all three levels)		radiatus	precipitatio	
	(none)	(none)	virga	
Cirrus	fibrosus	astoratus		
	uncinus	radiatus		
	spinosus	verrucatus	mamma	
	castellatus	duplicatus		
	floccus			
Cirrocumulus	stratiformis	undulatus	virga	
	lenticularis	radiatus		
	floccus	lacunosus	mamma	
Cirrostratus	fibrosus	duplicatus	(none)	
	nebulosus	undulatus		

HOW TO SPOT
CUMULUS CLOUDS

Cumulus are low, detached, puffy clouds that develop vertically in rising mounds, domes or towers, and have generally flat bases. Their upper parts often resemble cauliflower and they appear brilliant white when reflecting high sunlight, but can look dark when the sun is behind them. Cumulus tend to be randomly scattered across the sky.

TYPICAL ALTITUDES*:
2,000-3,000ft
WHERE THEY FORM:
Worldwide, except in Antarctica (the ground is too cold for thermals).
PRECIPITATION (REACHING GROUND): Generally none, except for brief showers from congestus.



Cumulus humilis




Cumulus mediocris



Cumulus congestus

CUMULUS SPECIES:
HUMILIS: Minimal vertical extent. They look flattened and appear wider than they are tall. Do not cause precipitation.
MEDIOCRIS: Moderate vertical extent. Might show protuberances and sproutings at the top. Appear as tall as they are wide. Do not cause precipitation.
CONGESTUS: Maximum vertical extent. The tops are like cauliflowers. Appear taller than they are wide. Cause brief downpours.
FRAGTUS: Ragged edges and broken up. Can form in the moist air below rain clouds.

CUMULUS VARIETIES:
RADIATUS: When Cumulus have formed into rows, or 'cloud streets', which are roughly parallel to the wind direction. Due to perspective, the rows appear to converge towards the horizon.



Cumulus mediocris radiatus

NOT TO BE CONFUSED WITH:
STRATOCUMULUS: Cumulus clouds are detached, not joined into a layer like Stratocumulus.
ALTOCUMULUS: Cumulus are not usually as regularly spaced as a layer of the higher AltoCumulus. The clouds also look larger than the clumps of the AltoCumulus. When they are above the cloudspotters, Cumulus appear larger than the width of three fingers, held at arm's length.
CUMULONIMBUS: which often develops from a large Cumulus congestus. A cloud is still a Cumulus when its upper region has a sharp outline, compared with the softer top of the Cumulonimbus.

* These approximate altitudes (above the surface) are for mid-latitude regions.

Top right: Michael Fisher (retrieved 19th November 2016); Bottom: Peter Cooper (retrieved 18/11/16)

HOW TO SPOT CUMULONIMBUS CLOUDS

Cumulonimbus are thunderstorm clouds, characterised by their enormous height. They are typically tall enough to reach the top of the troposphere, where they spread out in plumes of ice particles that can appear smooth, fibrous or striated. They have dark bases and produce heavy showers – often of hail – which can be accompanied by thunder and lightning.

TYPICAL ALTITUDES*:
2,000–45,000ft
WHERE THEY FORM:
Common in tropical and temperate regions. Rare in polar ones.
PRECIPITATION (REACHING GROUND):
Heavy downpours, often of hail.



Cumulonimbus calvus (mass 'bald')



Cumulonimbus capillatus (mass 'hairy')

CUMULONIMBUS SPECIES

The two species are distinguished by the appearance of the cloud's top.
calvus: When the upper region is of soft undines flattened mounds, without any fibrous or striated appearance.
capillatus: When the upper region is cirrus-like and fibrous or striated, often in the shape of an awl, plume or a disorderly mass of white hair.
CUMULONIMBUS VARIETIES:
There are no official varieties.

NOT TO BE CONFUSED WITH...
nimbostratus: which is a dark, ragged precipitating layer, covering the sky. It can look similar to a Cumulonimbus that is directly overhead (and also appears to cover much of the sky) but the precipitation will tend to be more steady and more persistent than the short heavy showers of the Cumulonimbus. If thunder, lightning or hail is present, then the cloud is a Cumulonimbus.
cumulus congestus: from which a Cumulonimbus often develops. Seen from a distance, the cloud is said to have changed into a Cumulonimbus when parts of its upper region begin to lose their sharp edges, due to the droplets freezing into ice crystals. Thunder, lightning or hail will also identify the Cumulonimbus.



* These approximate altitudes (above the surface) are for mid-latitude regions.

Top left: Bob Langford/Corbis; Top right: Alan Dyer/Corbis; Left: Bob Langford/Corbis; Right: Alan Dyer/Corbis

t
l
a
t
e

t
s
o
n
c
f
r
s
o
t

V
a
l
a
c
e
a
n
b
c
h

HOW TO SPOT STRATUS CLOUDS

Stratus are grey layers or patches of cloud, with very diffuse edges. They are the lowest-forming of all the cloud genera, sometimes appearing at ground level, when they are called fog or mist.

TYPICAL ALTITUDES*: 0–4,500ft
WHERE THEY FORM:
Worldwide. Most commonly around coasts and estuaries.
PRECIPITATION (REACHING GROUND):
No more than occasional drizzle, snow or snow grains.



Stratus fractus



Stratus fractus

SPECIES SPECIES:
None. Stratus is by far the most common, when it is in a grey, generally featureless layer.
fractus: When it is so opaque, ragged strands of grey cloud. This can appear in the region below precipitating clouds, when it is called 'lowered'. Though not particularly thick, these strands can look quite dark against the base of the cloud above.

STRATUS VARIETIES:
opacus: When the layer is thick enough to completely mask the sun or moon.
translucens: When it is thin enough to show the outline of the sun or moon.
conspicua: A rare variety, in which the layer has wave-like undulations to its surface. The surface of Stratus is rarely distinct enough for this to be observed.

NOT TO BE CONFUSED WITH...
cirrostratus: which is a high layer cloud that can look similar to a very thin Stratus. Being made of ice, however, it has a white tone.
altostratus: which is a mid-level layer cloud, often consisting of streaks, like Stratus. Though a layer of Stratus, the outline of the Sun (when it is discernible) is less diffuse, compared with the 'grounded' appearance of the Altostratus.
nimbostratus: which is a thick, dark layer of precipitating cloud that might be confused with a thick Stratus. But this has a less ragged base than the Nimbostratus and produces lighter precipitation.

* These approximate altitudes (above the surface) are for mid-latitude regions.

P
f

A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y
Z

**HOW TO SPOT
STRATOCUMULUS CLOUDS**

Stratocumulus are low layers or patches of cloud, with well-defined bases. They are usually composed of clumps or rolls, and often show strong variations in tone - from bright white to dark grey. Their cloud elements may be joined into continuous, unbroken layers or have gaps between them.

TYPICAL ALTITUDES*:
2,000-6,500ft
WHERE THEY FORM:
Worldwide - it's a very common cloud.
PRECIPITATION (REACHING GROUND):
Occasionally light rain, snow or snow pellets.




Stratocumulus stratiformis opacus... and *perlucidus*

STRATOCUMULUS SPECIES:
STRATIFORMIS: The most common, when the clumps or rolls extend over a large area. A 'roll cloud' is a particular formation, in the shape of a large, individual tube of cloud.
LENTICULARIS: When one or more mass of cloud is in a smooth, solid-looking almond or lens shape.
CASTELLANUS: When the elements have crenellated tops.

STRATOCUMULUS VARIETIES:
OPACUS: When the layer is thick enough to completely mask the sun or moon.
TRANSLUCIDUS: When it is thin enough to show the outline of the sun or moon.
PERLUCIDUS: When there are gaps between the cloud elements.
DUPLEXATUS: When there are layers at different altitudes, sometimes partly merged.
UNDULATUS: When the elements are arranged in nearly parallel lines.
KARMAIUS: When lines of closely bunched elements appear to converge towards the horizon.
LACINIOSUS: When the layer shows large net-like holes fringed with cloud.

NOT TO BE CONFUSED WITH:
CUMULUS: which is also clumpy, well defined, and forms at similar altitudes. The elements of Stratocumulus tend to be closer together and to have flatter tops.
ALTOCUMULUS: which is a mid-level layer of cloudlets. These appear smaller than the Stratocumulus elements, which - looking above 30° from the horizon - appear larger than the width of three fingers, held at arm's length.
STRATUS: which is a low, indistinct layer, with much less variation in tone and less definition than Stratocumulus.

* These approximate altitudes (above the surface) are for mid-latitude regions.

on
tog
aw
po
rar
fu
ab
est
th

cl
es
Ni
fer
co
so

pe
th
dr
wi
w
G
A

d
uj

S
b
cl
ai
g
it
te

**HOW TO SPOT
ALTOCUMULUS CLOUDS**


Alto cumulus are mid-level layers or patches of cloudlets, in the shape of rounded clumps, rolls or almonds/lenses. These are white or grey, and the sides away from the Sun are shaded. Alto cumulus are usually composed of droplets, but may also contain ice crystals.

TYPICAL ALTITUDES*:
6,500-18,000ft
WHERE THEY FORM:
Worldwide
PRECIPITATION (REACHING GROUND):
Very occasionally causes light rain.

ALTOCUMULUS SPECIES:
STRATIFORMIS: Most common, when the cloudlets extend over a large area.
LENTICULARIS: When it is in the form of one or more individual almond- or lens-shaped masses that appear dense, with pronounced shading.
CASTELLANUS: When the cloudlets have crenellated tops.
FLOCCUS: When the cloudlets are Cumulus-like tufts, with ragged bases, often with fibrous trails (virga) of ice crystals falling below.



Alto cumulus stratiformis undulans




Alto cumulus lenticularis *Alto cumulus floccus*

ALTOCUMULUS VARIETIES:
OPACUS: When the layer is thick enough to completely mask the sun or moon.
TRANSLUCIDUS: When it is thin enough to show the outline of the sun or moon.
PERALPICIDUS: When there are gaps between the cloudlets.
DUPLEXATUS: When there are layers at different altitudes, sometimes partly merged.
UNDULATUS: When the cloudlets are arranged in nearly parallel lines.
RADIATUS: When long lines of them appear to converge towards the horizon.
LACUNOSUS: When the layer shows net-like holes fringed with cloud.

NOT TO BE CONFUSED WITH...
CIRROCUMULUS: which is a higher layer of cloudlets, that appear like little grains of salt. Looking above 30° from the horizon, the larger Alto cumulus cloudlets generally appear the width of between one and three fingers, held at arm's length. Also, these exhibit shading, which those of Cirrocumulus don't.
CIRRUS: which is a high cloud, whose streaks of falling ice crystals can resemble Alto cumulus cloudlets showing virga, but do not have their dense-looking heads.

* These approximate altitudes (above the surface) are for mid-latitude regions.

http://www.met.rdg.ac.uk/~atm/122/122.htm
 http://www.met.rdg.ac.uk/~atm/122/122.htm

la
w
b
c
d
t
n
o
f
y
h
t


C
r
v
c
c
t
i
t
l

**HOW TO SPOT
ALTOSTRATUS CLOUDS**

Altostratus are mid-level layers of grey cloud, which are either featureless or fibrous in appearance, and typically extend over an area of several thousand square miles. Usually composed of both water droplets and ice crystals, they are often thin enough in parts to reveal the position of the sun, which appears as if through ground glass. Altostratus can cause a white or (when very thin) coloured 'cosoma' (disc of light) around the sun or moon.



Altostratus translucidus



Altostratus radiatus

<p>ALTOSTRATUS VARIETIES:</p> <p>OPACUS: When the cloud layer is generally thick enough to mask the position of the sun or moon.</p> <p>TRANSLUCIDUS: When it is generally thin enough to show the position of the sun or moon.</p> <p>DUPLEXATUS: When there is more than one layer at different altitudes, these often being partly merged. This is generally only visible when, by the light of a low sun, the higher layer is lit and the lower is in shadow, or when shearing winds cause the strata of the layers to drift.</p> <p>UNDULATUS: When the layer shows largely parallel undulations.</p> <p>RADIATUS: When lengthy undulations appear to converge toward the horizon.</p>	<p>NOT TO BE CONFUSED WITH...</p> <p>CIRROSTRATUS: which is a higher layer of ice crystals that looks like a thin, milky veil across the sky, and often thickens and lowers to develop into Altostratus. The Altostratus will tend to be more opaque, making the sunlight too diffuse for objects to cast shadows, as they do below Cirrostratus. White coloured or white discs of light, called 'cosmas', can appear around the sun/moon through Altostratus, this cloud will not cause the 'halo phenomena' of the Cirrostratus.</p> <p>NIMBOSTRATUS: which is a thick, dark layer of precipitating cloud that often develops out of an Altostratus. Generally darker, it produces considerably heavier rain or snow.</p>
---	--

* These approximate altitudes (above the surface) are for mid-latitude regions.

196. Peter Zetterstrom (November 1911), Ralph Albany Salsbery (November 1979)

t
i
c
J
t
c
t
C
s
tl
H

HOW TO SPOT
NIMBOSTRATUS CLOUDS

Nimbostratus are thick, grey, featureless layers of cloud that cause prolonged, continuous, often heavy, rain, snow or ice pellets. They tend to have very diffuse bases, as a result of all the falling precipitation. Nimbostratus are the deepest of all the layer clouds - sometimes extending from 2,000ft up to around 18,000ft - and generally extend over many thousand square miles. As with other precipitating clouds, the falling precipitation can cause Stratus fractus to form in the air below Nimbostratus clouds. These are known as 'pannus' and appear as shreds of cloud, looking darker than the underside of the Nimbostratus. When these join together, they tend to lower the bases of Nimbostratus clouds even further. They are invariably thick enough to completely hide the sun or moon.

TYPICAL ALTITUDES*:
2,000-18,000ft
WHERE THEY FORM:
Worldwide. More common in middle latitudes.
PRECIPITATION (REACHING GROUND): Causes moderate to heavy rain or snow (stray and prolonged).

NIMBOSTRATUS SPECIES:
There are no species, as the cloud's appearance is so uniform.

NIMBOSTRATUS VARIETIES:
There are no varieties, as the cloud's appearance is so uniform.

NOT TO BE CONFUSED WITH...
ALTOSTRATUS: which is a thinner - though also indistinct - layer of cloud. Nimbostratus is always darker than it and, by definition, produces precipitation. Altostratus only does sometimes, and this will generally be light. Whilst the position of

the sun can generally be determined through at least part of a layer of Altostratus, it will never be so through a Nimbostratus.
CUMulonimbus: which, when observed from directly below, can also appear as a very dark layer, covering the whole sky. The precipitation falling from a Nimbostratus will not generally be as heavy and will be more prolonged and continuous, compared with the sudden showers of the Cumulonimbus. Nor will the Nimbostratus produce its hail, thunder or lightning.



Nimbostratus - never a pretty sight

* These approximate altitudes (above the surface) are for mid-latitude regions.

David Brown (numbers 1127)

HOW TO SPOT
CIRRUS CLOUDS

Cirrus are the highest of the ten main cloud types. In the form of delicate, white streaks, patches or bands of falling ice crystals, they are detached from each other, and have fibrous or silky appearances. Cirrus rarely appear very thick. They are often seen with the other high clouds, Cirrostratus and Cirrocumulus and, like them, can show 'halo phenomena' around the sun or moon.



Cirrus uncinus

TYPICAL ALTITUDES*:
16,500-45,000ft
WHERE THEY FORM:
Worldwide
PRECIPITATION (REACHING GROUND): None



Cirrus floccus



Cirrus vertebratus

CIRRUS SPECIES:
FIBRATUS: When it is in the form of straight or curved filaments that are mostly distinct from each other and do not terminate in hooks or clumps.
UNCINUS: When its 'fallstreaks' are the shape of hooks or commas.
SPISSATUS: The thickest Cirrus - when it is in patches that appear grey in front of the sun - which tends to originate from the arrival of a Cumulonimbus.
CASTELLANUS: When it is in the form of small distinct clumps with crenellated tops.
FLOCCUS: When it is in the form of independent small round tufts, which often show trails of ice crystals falling from them.

CIRRUS VARIETIES:
INTORTUS: When the fallstreaks are irregular and tangled.
RADIATUS: When the filaments are in parallel bands, usually aligned to the wind at high altitude, which converge towards the horizon, due to perspective.

NOT TO BE CONFUSED WITH...
CIRROSTRATUS: which looks like a thin, milky smooth or fibrous veil across the sky. Cirrus, by contrast, is in separated streaks, fibres or patches.
CIRROCUMULUS: which is a high layer of cloudlets, like grains of salt. Cirrus does not show this finely dappled texture.

VERTEBRATUS: When the filaments look like a fish skeleton.
DUPLEXATUS: When the filaments, streaks or hooks are arranged at more than one altitude, which can be apparent when the winds cause them to point in different directions.

* These approximate altitudes (above the surface) are for mid-latitude regions.

Left: Malcolm Black (London, UK); Top right: Graham Thomas (Leeds, UK); Bottom right: (London, UK)

re
w
th
cl
'p
N
cr
sp
cl
th
di
of
Fc
pl
cl
re
di
th
co
tr
to
va
th
od

HOW TO SPOT
CIRROCUMULUS CLOUDS

Cirrocumulus are high patches of cloud or layers of tiny cloudlets that appear as white grains. These show no shading, even on the sides away from the sun. These cloudlets are generally regularly spaced, and often arranged in ripples, known as the undulatus variety.

TYPICAL ALTITUDES*:
16,500-45,000ft
WHERE THEY FORM:
Worldwide
PRECIPITATION
(REACHING GROUND):
None.

CIRROCUMULUS SPECIES:

STRATIFORMIS: When it is in an extensive layer, rather than just a patch. A less common species than for other genera.


LENTICULARIS: When it is in the form of one or more independent, well-defined, almond- or lens-shaped masses, which have smooth surfaces and are much larger than the grain-like cloudlets of the other species.

CASTELLANUS: When, on careful inspection, its cloudlets have castellated tops.

FLOCCUS: When, on careful inspection, its cloudlets are Cumulus-like, with ragged bases.



Cirrocumulus stratiformis



Cirrocumulus lacunosus undulatus

CIRROCUMULUS VARIETIES:

UNDULATUS: When its cloudlets are in a wave-like arrangement of ripples or broad undulations (or both at the same time).

LACUNOSUS: When the layer has holes fringed with cloud, like a net or honeycomb.

NOT TO BE CONFUSED WITH:

CIRRUS AND CIRROSTRATUS: which are streaks and smooth/fibrous layers of high cloud. Cirrocumulus layers, by contrast, are subdivided into many grain-like cloudlets.

ALTOCUMULUS: which is a mid-level layer of larger cloudlets. Looking above 30° from the horizon, the smaller Cirrocumulus cloudlets generally appear less than the width of one finger, held at arm's length.

* These approximate altitudes (above the surface) are for mid-latitude regions.

© 2000 Sky & Telescope Magazine

C
Inde
a tra
the :
way
appe
reco
delig
T
level
is nc
how
sky
varit
strat
a te
that
is ps
shap

THI
high
as t
rese
app
A
dete
ther
tem
depi
form
imp
surf
the

